

Kinetic non-optimality and vibrational stability of proteins

Marek Cieplak* and Trinh Xuan Hoang

Institute of Physics, Polish Academy of Sciences, Al. Lotnikow 32/46, 02-668 Warsaw, Poland

*Correspondence to:

Marek Cieplak,

Physics Department,

Rutgers University,

Piscataway NJ-08854, USA.

Tel: 732-445-1440

Fax: 732-445-4343

E-mail: cieplak@physics.rutgers.edu

Grant sponsor: KBN (Poland); Grant number: 2P03B-146-18.

Keywords: protein folding; Go model; molecular dynamics; protein stability

Abstract

Scaling of folding times in Go models of proteins and of decoy structures with the Lennard-Jones potentials in the native contacts reveal power law trends when studied under optimal folding conditions. The power law exponent depends on the type of native geometry. Its value indicates lack of kinetic optimality in the model proteins. In proteins, mechanical and thermodynamic stabilities are correlated.

INTRODUCTION

Proteins are extraordinary heteropolymers. They fold to their native states much faster than what a blind combinatorics would predict¹ since a folding funnel in the energy landscape is formed.²⁻⁴ Proteins are believed to have high designabilities,⁵ to be stable against mutations,^{6,7} and to have the highest densities of states.⁸ Furthermore, the α helix secondary motifs have been shown theoretically to be the fastest folders among chains of the same number, N , of aminoacids⁹ and be the result of the geometrical optimization of compact chains with maximum wiggle room.¹⁰ Experimental results¹¹⁻¹³ (see also a commentary by Chan¹⁴) also point to the accelerating role of the helices. Biological evolution may have optimized functionality of proteins, but can proteins be optimal kinetically?

Here, we consider folding times, t_{fold} , in Go models¹⁵⁻¹⁷ of proteins and decoy structures and show that though proteins fold to their native structures fast they are not optimal folders. This conclusion ties well with the protein engineering experiments^{18,19} which show that mutations in wild type proteins may lead to significant increases in folding rates and thus show no kinetic optimality of sequences. Our theoretical argument is based on relating universality classes in the scaling of t_{fold} to classes of native geometries. This confirms a decisive role of native geometry in determining properties of proteins.²⁰ The scaling trends

that we observe are robust when studied at the temperature of the fastest folding, T_{min} , but become obscure when studied at other temperatures.

Another issue which we examine here regards the notion of protein stability. One definition of stability is thermodynamic – it assesses the role of non-native phase space valleys relative to the native valley by determining the probability of staying in the native basin. It is characterized by the folding temperature, T_f , at which this probability crosses $\frac{1}{2}$.²¹ Another is mechanical: at what temperature will the native conformation melt due to vibrations? The mechanical definition does not refer to non-native valleys. The two notions should correlate with each other if the native valley dominates in the energy landscape. We show that this is indeed what happens in model proteins.

MODEL AND METHOD

We first consider the problem of universality classes in the scaling of folding properties. There have been various predictions about the nature of scaling of t_{fold} . A number of theories suggested a power law dependence of barrier heights on N and thus an exponential law for t_{fold} .^{22–24} Thirumalai,²⁵ however, has argued in favor of a power law for t_{fold} ,

$$t_{fold} \sim N^\lambda, \quad (1)$$

where λ is estimated to be between 3.8 and 4.2 for simple two-state folders. A heuristic model²⁶ leads to $\lambda = 3$. Numerical studies of t_{fold} in various lattice models^{27–29} have supported the power law behavior and indicated dependence of λ on specifics of the model, dimensionality and temperature, T . For designed sequences in three dimensions, λ has been found to be in the Thirumalai range²⁷ whereas for Go models it has been found to be of order 3.^{27,29} In these studies, t_{fold} is defined as the first passage time.

Here, we extend the scaling studies to off-lattice Go models¹⁵ and consider chains of beads separated by $d_0 \approx 3.8\text{\AA}$ – a typical length of the peptide bond. The Go Hamiltonian is defined through a native conformation of a sequence since it assigns relevant interaction

energies only to the native contacts. Despite this simplification, the Go models may behave more realistically than atomistic models.³⁰

It should be noted that the Go models are so minimal that they disregard an explicit amino acidic definition of a protein and variability of the volume taken by individual side chains. Natural proteins appear to fold by locking its segments together in an unfrustrated way. Adding attraction to the non-native contacts in the bead-spring model might seem to be making the model more realistic but, in fact, it leads to spurious entanglements during folding. In this sense, the Go model repairs some of its shortcomings by a mutual cancellation of its ills and focuses on the effects related to the native structure. This focus is justified by experimental indications that the native structure itself is central to folding.^{31,13} On the other hand, the target oriented aspects of such theoretical modeling are hard to justify on a fundamental level.³² The nature of the Go model allows one to study the role of the native structure in kinetics but it does not allow to address the role of the sequential order. Determining sequence based, as opposed to structure based, classes of kinetic universality would be much more interesting but, clearly, also much more challenging.

We employ Lennard-Jones (LJ) potentials $V_{ij} = 4\epsilon \left[\left(\frac{\sigma_{ij}}{r_{ij}} \right)^{12} - \left(\frac{\sigma_{ij}}{r_{ij}} \right)^6 \right]$ for the native contact interactions between monomers i and j , in a distance of r_{ij} apart. The non-native interactions are described by repulsive soft core potentials that provide excluded volume and prevent entanglements. Our approach is presented in details in Ref.^{16,17} where several secondary structures and three model proteins have been analyzed. Such models were also studied in Ref.^{33,34} The distances between successive beads are controlled by an anharmonic potential. The length parameters σ_{ij} are selected so that the minimum of V_{ij} corresponds to the geometry found in the target structure and the contacts are said to be formed when i and j are not consecutive along the chain and r_{ij} is less than d_{nat} , where d_{nat} is 7.5Å.

There are other variants of the off-lattice Go models: Zhou and Karplus³⁵ and Dokholyan et al.³⁶ have considered models with a square well potential. Clementi et al.³⁷ have studied the 12-10 power law potentials. It is not clear which effective potential is the best and our choice is LJ.

The dynamics of the system are described by the Langevin equation $m\ddot{\mathbf{r}} = -\gamma\dot{\mathbf{r}} + F_c + \Gamma$ where $\mathbf{r}$ is a position of a monomer, m is its mass and F_c is the force derived from the Hamiltonian. γ is a friction coefficient and Γ is the random force such that $\langle \Gamma(0)\Gamma(t) \rangle = 2\gamma k_B T \delta(t)$, where k_B is the Boltzmann constant, t is time and $\delta(t)$ is the Dirac delta function. Both the friction and the random force represent the effects of the solvent and they control T . The equations are solved using the fifth order predictor-corrector scheme.

In the following, T is measured in the units of ϵ/k_B and t is measured in units of the oscillatory period τ . At low values of friction, τ is equal to $\sqrt{ma^2/\epsilon}$, where a is a van der Waals radius of the amino acid residues. The value of a is chosen as $5\text{\AA}$ which is roughly equal to $\langle \sigma_{ij} \rangle$ in our model proteins. The simulations are done with $\gamma = 2m\tau^{-1}$ – a standard choice in studies of liquids. Higher values of γ have been argued to be more realistic.³⁸ We have shown¹⁶ that t_{fold} is linear in γ and T_{min} depends on γ weakly.

The native conformation is defined through the locations of the α carbons. We have considered 21 single domain Protein Data Bank (PDB) structures³⁹ with N ranging between 29 and 98. 9 of these structures belong to a set of proteins considered by Plaxco et al.¹² or are their close homologues. These are: the SH3 domain of 1efn (57), 2ptl (63), 2ci2 (83-18=65; 18 are not resolved), 1csp (67), 1ubq (76), 1hdn (85), 2abd (86), 1ten (90), and 1aps (98), where the numbers in brackets indicate the corresponding values of N . The additional 12 structures are: 1cti (29), 1cmr (31), 1ce4 (35), 1bba(36), 1erc (40), 1crn (46), 7rxn (52), 5pti (58), 1tap(60), 1aho (64), 1ptx (64), 1erg (70). These conformations were picked from the low- N end of the size distribution to allow for a reliable characterization. Our studies of these structures indicate well defined overall trends in t_{fold} which are only weakly affected by an inclusion of steric constraints.³⁸ Our results will be given here only for models without such constraints.

The results obtained for the PDB structures are compared to five classes of decoy conformations which differ in the way they fill space and in their packing arrangements. These classes form statistical ensembles in which a given value of N has multiple realizations. Four classes are defined in terms of shapes that homopolymers arrive at under various cooling

procedures. The non-consecutive beads in the homopolymers interact through the LJ potential with $\sigma_{ij} = 5\text{\AA}$ which corresponds to a typical van der Waals radius of aminoacids. We discuss the following classes (see Figure 1):

HC: conformations obtained through slow homopolymer cooling. The procedure involves generating an open conformation, assigning identical strengths to all inter-bead interactions, and then slowly annealing. The resulting compact conformation serves as a native structure in the Go-like Hamiltonian.

HQ: similar to HC but with a rapid quenching instead of annealing. The procedure results in non-compact native structures which, however, have many local contacts, as measured along the chain, and are thus more closely related to α -helices than to random heteropolymers.

HA: similar to HC but the α -helices of various lengths (of order 15) are first built into the initial states in a way which is consistent with the LJ couplings and then preserved through the annealing process by assigning ten times stronger couplings to the helical secondary structures.

HB: similar to HA but the helical segments are replaced by β -sheet conformations. The lengths of the β -strands are fixed at 8 monomers.

CL: compact native conformations generated on a grid as a self-avoiding random walk within a compact box of lattice constant equal to the length of a peptide bond and then stabilized by appropriate Lennard-Jones interactions.

The folding properties are studied as a function of T and then presented here for $T=T_{min}$ and T_f . at which probability, P_0 , of being in the native basin is $\frac{1}{2}$. P_0 is determined based on 10-15 long molecular dynamics trajectories at equilibrium. The results are illustrated in Figure 2 for two model proteins 1ubq and 1ce4.

The median folding times depend on T in a U-shaped fashion and, generally, the bigger the N , the narrower the U. The dependence of t_{fold} for the Go models of 1ubq and 1ce4 is shown at the bottom of Figure 2.

The system is assumed to be in its native state if all of its native contacts are established. A native contact between monomers i and j is said to be established if r_{ij} is shorter than $1.5\sigma_{ij}$.

RESULTS AND DISCUSSION

Kinetic Universality Classes

Figure 3 shows the validity of the power law, eq. (1), for t_{fold} when determined under the most favorable kinetic conditions – at T_{min} . The exponent λ sensitively depends on the geometrical class of the native structures (Table I). The case of HB is special since a crossover between two effective values of λ is observed (the β -strand lengths of 8 impose a condition on N above which a characteristic β -sheet behavior can start to be seen). The values of λ range between 1.7 and 3.2. The smallest λ corresponds to the HA and the largest to the HQ and long HB conformations. HC is intermediate. Note that λ for HA is smaller than 2 – the value suggested by de Gennes⁴⁰ in his analysis of the time scale for the coil to globule transition of a homopolymer.

The data points for PDB at T_{min} are somewhat scattered – there is no averaging over an ensemble – but a well defined trend is visible. The exponent λ is about 2.5 ± 0.2 which indicates that these structures are not optimal kinetically. HA, short HB, and HC of the same N fold faster. PDB appears to be comparable to the grid conformations CL (there is only a weak dependence on the dimensionality in the off-lattice models – when the grid structures are generated on the square lattice, λ becomes equal to 2.1 ± 0.2).

The existence of a trend in the scaling of t_{fold} for the PDB structures appears to be at odds with the analysis of experimental data compiled by Plaxco et al.,¹² and replotted here

in Figure 4, which indicated lack of any correlations with N . A flavor of this is already seen in Figure 3 which shows that one sequence (1aps) has a t_{fold} which is distant from the trend. This sequence appears to be frustrated geometrically and it has a very small experimental folding rate¹² – maybe it is just a poor folder. However, the experimental data indicate a significantly larger scatter of values than as seen in Figure 3.

There are three explanations of the discrepancy that we considered. First: the range of the values of N considered in Ref.¹² is smaller than studied here, which in itself emphasizes fluctuations. However, in data that were published later¹³ the range of N was extended to about 150 and the correlations of kinetics with N remained weak so the limited range of the values of N is not a likely explanation of the discrepancy. Second, it is only the simplified models, like the Go models, that show trends in the kinetics of folding whereas any additional complexities present in real systems may perturb such trends beyond detection. This possibility could be studied in the future by considering scaling in more complicated classes of models. In particular, the role of the localization index of the interactions should be elucidated in the context of scaling. Third: the trends are derailed by the fact that the experimental data are usually obtained at a fixed temperature, typically, but not necessarily, at the room temperature. Thus the data collection involved no kinetic optimization which would require selecting the best T for each protein individually.

The role of this third possibility is illustrated at the top of Figure 4 which shows the scaling of t_{fold} at T_f . The scatter is seen to be significantly larger than at T_{min} . It is not as large as in the experimental data but it should be noticed, again, that our Go systems are just very simple models of systems which are quite complicated. Another way to assess the relevance of the optimal selection of T is shown in Figure 5 which reanalyzes the data of figures 3 and 4 so that theoretically determined t_{fold} is plotted against the experimentally measured folding time, t_{exp} . The small number of available points makes it hard to place a bet on the best trend. However, it is clear that the points determined at T_{min} exhibit significantly less scatter than those calculated at T_f . This finding gives further support for the idea that the lack of optimization in the temperature may mask existence of any

underlying trends.

Our findings on scaling of characteristic T 's can be summarized as follows. For HC, HA and CL, T_{min} grows with N whereas T_f is almost constant and somewhat lower than T_{min} . The difference between T_{min} and T_f grows most slowly for HA. For PDB, T_{min} does not seem to have a trend, within the range of N studied, and the values of T_{min} are usually just above the corresponding values of T_f . This indicates a borderline behavior between excellent and poor folding characteristics, if the condition for the latter is $T_f \ll T_{min}$.^{21,41} This borderline behavior might characterize classes of proteins, especially of those which have a short lifetime in a living cell but this result may depend on the choice of the potentials.

Stability Against Vibrations

We now discuss stability of the native state in proteins. The mechanical stability can be probed through the phononic spectra as in Ref.¹⁶. This is accomplished by determining the frequency gap, ω_1 in the low end of the frequency spectrum. Another test is provided by studying root mean square displacements around the native state and employing the Lindemann criterion for melting. We introduce the parameter

$$\delta_L(T) = \frac{1}{n} \sum_{i>j}^{[NAT]} \frac{\sqrt{\langle r_{ij}^2 \rangle - \langle r_{ij} \rangle^2}}{r_{ij,NAT}}, \quad (2)$$

which is a variant of the parameter used by Takano et al.⁴² The summation is over pairs of monomers (n of them) which form native contacts and their rms displacement is compared to the native distances. The temperature, T_L , at which δ_L crosses the Lindemann value of 0.1 is a measure of mechanical stability. Figure 6 shows that both ω_1 and T_L show a correlation with T_f which suggests the predominance of the native valley in the energy landscape. Note that T_L is higher than T_f which indicates that the probability "leaks out" of the native state already when the vibrations in the native valley are small. Thus for good folders, the notions of thermodynamic and mechanical stabilities qualitatively coincide.

CONCLUSIONS

Our main results on scaling can be summarized as follows. There are kinetic universality classes among well folding sequences. These classes depend on the type of geometry involved in the native state. Well defined scaling trends can be established if folding is studied under optimal conditions. Otherwise they are hard to be seen, especially if the range of system sizes is narrow. The shapes of actual proteins in their native states are such that the folding times scale with an exponent which is higher than certain artificial classes of structures. This suggests the lack of kinetic optimality of proteins.

Our results have been obtained within the Go model which focuses on the role of the native state geometry. This level of simplified description incorporates a long list of approximations which somehow appear to compensate mutually. In effect, the Go systems are reasonable models of good folders and the simplifications involved are precisely of the kind that allow for a statistical analysis necessary to establish scaling properties. Working with sequences described in a more sophisticated manner would add to the reality of description. However, it would also necessitate dealing with statistical ensembles of sequences defined by more parameters than just the size and shape and currently that would be prohibitive numerically. Our results should then be viewed as establishing first inroads into understanding of the role of size in folding kinetics.

ACKNOWLEDGMENTS

We appreciate discussions with J. R. Banavar, C. Johnson, H. Nymeyer, J. N. Onuchic, S. Plotkin, and D. Thirumalai which motivated parts of this research. This work was funded by KBN.

REFERENCES

- ¹ Levinthal C. In: Debrunner P, editor. Mossbauer Spectroscopy in Biological Systems, Urbana: University of Illinois Press; 1969.
- ² Leopold PE, Montal M, Onuchic JN. Protein folding funnels: a kinetic approach to the sequence-structure relationship. *Proc Natl Acad Sci USA* 1992;89:8721-8725.
- ³ Bryngelson JD, Onuchic JN, Socci ND, Wolynes PG. Funnels, pathways, and the energy landscape of protein-folding – a synthesis. *Proteins: Struct Funct Genet* 1995;21:167-195.
- ⁴ Dill KA, Chan HS. From Levinthal to pathways to funnels. *Nature Struct Biol* 1997;4:10-19.
- ⁵ Li H, Tang C, Wingreen NS. Are protein folds atypical?. *Proc Natl Acad Sci* 1998;95:4987-4990.
- ⁶ Vendruscolo M, Maritan A, Banavar JR. Stability threshold as a selection principle for protein design. *Phys Rev Lett* 1997;78:3967-3970.
- ⁷ Bornberg-Bauer E, Chan HS. Modeling evolutionary landscapes: mutational stability, topology, and superfunnels in sequence space. *Proc Natl Acad Sci USA* 1999;96:10689-10694.
- ⁸ Micheletti C, Banavar JR, Maritan A, Seno F. Protein Structures and optimal folding from a geometrical variational principle. *Phys Rev Lett* 1999;82:3372-3375.
- ⁹ Maritan A, Micheletti C, Banavar JR. Role of secondary motifs in fast folding polymers: a dynamical variational principle. *Phys Rev Lett* 2000;84:3009-3012.
- ¹⁰ Maritan A., Micheletti C., Trovato A., Banavar JR. Optimal shapes of compact strings. *Nature* 2000;406:287-290.
- ¹¹ LopezHernandez E., Cronet P., Serrano L., Munoz V. Folding kinetics of Che Y mutants with enhanced native alpha-helix propensities. *J Mol Biol* 1997;266:610-620.

- ¹² Plaxco KW, Simons KT, Baker D. Contact order, transition state placement and the refolding rates of single domain proteins. *J Mol Biol* 1998;277:985-994.
- ¹³ Plaxco KW, Simons KT, Ruczinski I, Baker D. Topology, stability, sequence, and length: defining the determinants of two-state protein folding kinetics. *Biochemistry* 2000;39:11177-11183.
- ¹⁴ Chan HS. Matching speed and locality. *Nature* 1998;392:761-763.
- ¹⁵ Abe H, Go N. Noninteracting local-structure model of folding and unfolding transition in globular proteins. II. Application to two-dimensional lattice proteins. *Biopolymers* 1981;20:1013-1031.
- ¹⁶ Hoang TX, Cieplak M. Molecular dynamics of folding of secondary structures in Go-like models of proteins. *J Chem Phys* 2000;112:6851-6862.
- ¹⁷ Hoang TX, Cieplak M. Sequencing of folding events in Go-like proteins. *J. Chem. Phys.* 2001;113:8319-8328.
- ¹⁸ Kim DE, Gu H, Baker D. The sequences of small proteins are not extensively optimized for rapid folding by natural selection. *Proc Natl Acad Sci* 1998;95:4982-4986.
- ¹⁹ Fersht AR. Transition state structure as a unifying basis in protein folding mechanisms: contact order, chain topology, stability and the extended nucleus mechanism. *Proc Natl Acad Sci* 2000;97:1525-1529.
- ²⁰ Baker D. A surprising simplicity to protein folding. *Nature* 2000;405:39-42.
- ²¹ Socci ND, Onuchic JN. Folding kinetics of protein-like heteropolymers. *J Chem Phys* 1994;101:1519-1528.
- ²² Takada S, Wolynes PG. Microscopic theory of critical folding nuclei and reconfiguration activation barriers in folding proteins. *J Chem Phys* 1997;107:9585-9598.
- ²³ Finkelstein AV, Badredtinov AY. Rate of protein folding near the point of thermodynamic

- equilibrium between the coil and the most stable chain fold. *Fold Des* 1997;2:115-121.
- ²⁴ Wolynes PG. Folding funnels and energy landscapes of larger proteins within the capillarity approximation. *Proc Natl Acad Sci USA* 1997;94:6170-6175.
- ²⁵ Thirumalai D. From minimal models to real proteins: timescales for protein folding. *J Physique I* 1995;5:1457-1467. Thirumalai D, Klimov DK. Deciphering the timescales and mechanisms of protein folding using minimal off-lattice models. *Curr Opin Struct Biol* 1999;9:197-207.
- ²⁶ Camacho CJ. Entropic barriers, frustration, and order: basic ingredients in protein folding. *Phys Rev Lett* 1996;77:2324-2327.
- ²⁷ Gutin AM, Abkevich VI, Shakhnovich EI. Chain length scaling of protein folding time. *Phys Rev Lett* 1996;77:5433-5436.
- ²⁸ Zhdanov VP. Folding time of ideal β sheets vs. chain length. *Europhys Lett* 1998;42:577-581.
- ²⁹ Cieplak M, Hoang TX, Li MS. Scaling of folding properties in simple models of proteins. *Phys Rev Lett* 1999;83:1684-1687.
- ³⁰ Takada S. Go-ing for the prediction of protein folding mechanism. *Proc Natl Acad Sci USA* 1999;96:11698-11700.
- ³¹ Alm E., Baker D. Matching theory and experiment in protein folding. *Curr Opin Struct Biol* 1999;277:189-196.
- ³² Kaya H, Chan HS. Polymer principles of protein calorimetric two-state cooperativity. *Proteins: Struct Funct Genet* 2000;40:637-661.
- ³³ Li MS, Cieplak M. Folding in two-dimensional off-lattice models of proteins. *Phys Rev E* 1999;59:970-976.
- ³⁴ Klimov DK, Thirumalai D. Mechanisms and kinetics of β -hairpin formation. *Proc Natl*

Acad Sci USA 2000;97:2544-2549.

- ³⁵ Zhou Y, Karplus M. Interpreting the folding kinetics of helical proteins. *Nature* 1999;401:400-403.
- ³⁶ Dokholyan NV, Buldyrev SV, Stanley HE, Shakhnovich EI. Discrete molecular dynamics studies of the folding of a protein-like model. *Folding Des* 1998;3:577-587.
- ³⁷ Clementi C, Nymeyer H, Onuchic JN. Topological and energetic factors: what determine the structural details of the transition state ensemble and “on-route” intermediates for protein folding? An investigation for small globular proteins. *J Mol Biol* 2000;298:937-953.
- ³⁸ Veitshans T, Klimov D, Thirumalai D. Protein folding kinetics: time scales, pathways and energy landscapes in terms of sequence-dependent properties. *Folding Des* 1997;2:1-22.
- ³⁹ Bernstein FC, Koetzle TF, Williams GJB, Meyer Jr. EF, Brice MD, Rodgers JR, Kennard O, Shimanouchi T, Tasumi M. The Protein Data Bank: a computer-based archival file for macromolecular structures. *J Mol Biol* 1977;112:535-542.
- ⁴⁰ de Gennes PG. Kinetics of collapse for a flexible coil. *J Phys Lett* 1985;46:L639-L642.
- ⁴¹ Cieplak M, Henkel M, Karbowski J, Banavar JR. Master equation approach to protein folding and kinetic traps. *Phys Rev Lett* 1998;80:3654-3657.
- ⁴² Takano M, Takahashi T, Nagayama K. Helix-coil transition and $1/f$ fluctuation in a polypeptide. *Phys Rev Lett* 1998;80:5691-5694.

TABLES

TABLE I. The values of exponent λ for the classes of conformations studied.

Structure	λ
HC	2.2 ± 0.1
HA	1.7 ± 0.1
HB	$0.9 \pm 0.1, 3.2 \pm 0.1$
HQ	2.7 ± 0.2
CL	2.6 ± 0.2
PDB	2.5 ± 0.2

FIGURE CAPTIONS

Fig. 1. Examples of native conformations that were used in these studies. The folding data were generated based on 11 realizations of each class of structures for each value of N , except for the case of HA when 5 realizations were sufficient.

Fig. 2. Equilibrium and kinetic properties of the Lennard-Jones-Go model of two proteins, 1ce4 (solid lines) and 1ubq (dotted lines), as a function of temperature. The top panel shows the probability of staying in the native state and the bottom panel shows the median folding time as determined based on 200 trajectories at each temperature.

Fig. 3. Power law dependence of the median folding times at T_{min} on N for the indicated classes of geometry of the native conformations. The proteins analyzed by Plaxco et al.¹² are shown as open squares. Closed squares correspond to other proteins. For CL, the times are multiplied by 10.


Fig. 4. Top: Values of t_{fold} for the PDB structures as determined at T_f . The open circles refer to the proteins studied by Plaxco et al.¹² The line shows the scaling trend found at T_{min} . Bottom: Experimentally determined folding times, t_{exp} , (inverses of the folding rates) as compiled by Plaxco et al.¹²

Fig. 5 Folding times of the theoretical Go model versus folding times observed experimentally in proteins studied by Plaxco et al.¹². The solid and open squares correspond to $T = T_{min}$ and $T = T_f$ respectively.


Fig. 6. Bottom: T_L as a function of T_f for the PDB structures studied. Top: The lowest non-zero phononic frequency of the same structures plotted vs. T_f . The broken lines indicate overall trends. The inset illustrates the dependence of δ_L (eq. 2) on temperature, T , for the Go model of crambin. The horizontal line indicates the 10% value of δ_L .

FIGURES


PDB (1aho)




CL




HC




HQ



HA



HB



5Å

FIG. 1.




FIG. 2.




FIG. 3.




FIG. 4.




FIG. 5.




FIG. 6.